



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

general reader on the various phases of bird migration should be more or less vitiated throughout by the ill-devised theory which pervades and colors an otherwise praiseworthy book,—a work, in other respects, as regards its general character, far in advance of Mr. Dixon's previous one bearing the same title.—J. A. A.

**Marsh on the Affinities of *Hesperornis*.**<sup>1</sup>—Professor Marsh here reaffirms the correctness of his conclusion, published in 1880, that “the Struthious characters, seen in *Hesperornis*, should probably be regarded as evidence of real affinity, and in this case *Hesperornis* would be essentially a carnivorous, swimming Ostrich.” Authors who had not seen the original specimens, says Prof. Marsh, “seem to have accepted without hesitation the striking adaptive characters of the posterior limbs as the key to real affinities,” till soon “the Ratite affinities of *Hesperornis* were seldom alluded to in scientific literature.” He has remained silent, “leaving to future discoveries the final decision of the question at issue.” This decision, Prof. Marsh thinks, is now on record, Prof. Williston having discovered near the original type locality a remarkably perfect specimen of *Hesperornis*, with the feathers in place, showing that *Hesperornis* had “the typical plumage of an Ostrich.” Reference to Prof. Williston's paper (Kansas University Quarterly, Vol. V, No. 1, July, 1896, pp. 53, 54, pl. ii) shows that there is still ground for a difference of opinion as to the Struthious character of the downy feathers found on the tarsus and head of Prof. Williston's specimen of *Hesperornis*.—J. A. A.

**Stone on the Genus *Sturnella*.**<sup>2</sup>—Mr. Stone's paper has relation mainly to the forms referred to *S. magna mexicana*, the Rio Grande Valley phase of which group Mr. Stone now separates as a new subspecies, under the name *S. m. hoopesi*. This form resembles *magna* in the coloration of the lower parts, it lacking the yellow on the malar region, while the upper plumage is lighter even than in *neglecta*, with the tail bars “more distinct than in any of the other races.” True *S. m. mexicana* thus becomes restricted to southern Mexico and Central America, *S. m. hoopesi* taking its place in the A. O. U. Check-List. The Florida bird, which has sometimes been referred to *mexicana*, Mr. Stone finds is not separable from Louisiana examples, and that these latter differ but little from specimens from southern Indiana and southern Illinois. He considers it therefore inadvisable to separate this Gulf coast phase from *magna*.—J. A. A.

---

<sup>1</sup> The Affinities of *Hesperornis*. By O. C. Marsh. American Journal of Science, III, April, 1897, pp. 347, 348.

<sup>2</sup> The Genus *Sturnella*. By Witmer Stone. Proc. Acad. Nat. Sci. Phila., 1897, pp. 146–152.